

## Cuvitru

## **Prior Authorization Request**

CVS Caremark administers the prescription benefit plan for the patient identified. This patient's benefit plan requires prior authorization for certain medications in order for the drug to be covered. To make an appropriate determination, providing the most accurate diagnosis for the use of the prescribed medication is necessary. **Please respond below and fax this form to CVS Caremark toll-free at 1-855-330-1720**. If you have questions regarding the prior authorization, please contact CVS Caremark at **1-888-877-0518**. For inquiries or questions related to the patient's eligibility, drug copay or medication delivery; please contact the Specialty Customer Care Team: CaremarkConnect® 1-800-237-2767.

The recipient of this fax may make a request to opt-out of receiving telemarketing fax transmissions from CVS Caremark. There are numerous ways you may opt-out: The recipient may call the toll-free number at 877-265-2711, at any time, 24 hours a day/7 days a week. The recipient may also send an opt-out request via email to do not call@cvscaremark.com. An opt out request is only valid if it (1) identifies the number to which the request relates, and (2) if the person/entity making the request does not, subsequent to the request, provide express invitation or permission to CVS Caremark to send facsimile advertisements to such person/entity at that particular number. CVS Caremark is required by law to honor an opt-out request within thirty days of receipt.

Patient's Name:	Date:
Patient's ID:	Patient's Date of Birth:
Physician's Name:	
Specialty:	
Physician Office Telephone:	
<b>Referring</b> Provider Info: □ Same as Re	equesting Provider
Name:	NPI#:
Fax:	
	eferring Provider   Same as Requesting Provider
Name:	NPI#:
Fax:	Phone:
	t to dosing limits in accordance with FDA-approved labeling, pendia, and/or evidence-based practice guidelines.
Required Demographic Information:	
Patient Weight:	kg
Patient Height:	ст

	e of Service Questions:  Where will this drug be administered?  ☐ Ambulatory surgical, skip to Clinical Questions ☐ Off-campus Outpatient Hospital ☐ Physician office, skip to Clinical Questions	<ul> <li>☐ Home infusion, skip to Clinical Questions</li> <li>☐ On-campus Outpatient Hospital</li> <li>☐ Pharmacy, skip to Clinical Questions</li> </ul>
В.	Is the patient less than 21 years old or 65 years of age or ☐ Yes − less than 21 years old, <i>skip to Clinical Criteria</i> ☐ Yes − age 65 years or older, <i>skip to Clinical Criteria</i> ☐ No	Questions
C.	Is this request to continue previously established treatment. ☐ Yes - This is a continuation of an existing treatment. ☐ No - This is a new therapy request (patient has not reconcilinated Criteria Questions	•
D.	Has the patient experienced an adverse event with the requinterventions (eg acetaminophen, steroids, diphenhydrami event (anaphylaxis, anaphylactoid reactions, myocardial in immediately after an infusion? <i>ACTION REQUIRED: A</i> ☐ Yes, <i>skip to Clinical Criteria Questions</i> ☐ No	ne, fluids or other pre- medications) or a severe adverse nfarction, thromboembolism, or seizures) during or
Е.	Does the patient have laboratory confirmed anti-IgA anti-clinical documentation.   Yes, skip to Clinical Criteria	
F.	Is the patient medically unstable which may include respir the member's ability to tolerate a large volume or load or cannot be managed in an alternate setting without appropr REQUIRED: Attach supporting clinical documentation	predispose the member to a severe adverse event that iate medical personnel and equipment? <i>ACTION</i>
G.	Does the patient have significant behavioral issues and/or safety of the infusion therapy AND the patient does not has <i>supporting clinical documentation</i> .	

Cr	iteria Questions:	
1.	What is the diagnosis? List continues on following page.	
	☐ Myasthenia gravis	☐ Macrophage Activation Syndrome (MAS
	☐ Chronic inflammatory demyelinating polyneuropathy (CIDP)	☐ Hyperimmunoglobulinemia E syndrome
	☐ Immune thrombocytopenic purpura (ITP)	☐ Multiple myeloma
	☐ B-cell chronic lymphocytic leukemia (CLL)	☐ Opsoclonus-myoclonus
	☐ Stiff-person syndrome	☐ Post-transfusion purpura
	☐ Dermatomyositis	☐ Solid organ transplantation
	□ Polymyositis	☐ Stevens-Johnson syndrome
	☐ Multifocal motor neuropathy	☐ Toxic necrotizing fasciitis
	☐ Human immunodeficiency virus (HIV) infection	☐ Toxic epidermal necrolysis
	☐ Guillain-Barré syndrome	☐ Toxic shock syndrome
	☐ Lambert-Eaton myasthenic syndrome	☐ Kawasaki syndrome (pediatric)
		☐ Isoimmune hemolytic disease of newborn
	Parvovirus B19-induced pure red cell aplasia	
	☐ Fetal/neonatal alloimmune thrombocytopenia	□ Neonatal hemochromatosis
	☐ Immune checkpoint inhibitor related toxicity	☐ Acquired red cell aplasia
	☐ CAR-T therapy related hypogammaglobulinemia	☐ Acute disseminated encephalomyelitis
	Rasmussen encephalitis	☐ Autoimmune neutropenia
	☐ Enteroviral meningoencephalitis	☐ Autoimmune hemolytic anemia
	☐ Systemic lupus erythematosus	☐ Autoimmune neutropenia
	☐ Hematophagocytic lymphohistiocytosis (HLH)	☐ BK virus associated nephropathy
	☐ Major surgery associated secondary immunosuppression	☐ Churg-Strauss syndrome
	☐ Major burns associated secondary immunosuppression	☐ Birdshot retinochoroidopathy
	☐ Hematologic malignancy associated secondary immunosuppress	ion
	☐ Collagen-vascular disease associated secondary immunosuppress	sion
	☐ Bone marrow transplant/hematopoietic stem cell transplant recip	
	☐ Autoimmune mucocutaneous blistering disease (includes pemph	
	pemphigoid, mucous membrane pemphigoid, and epidermolysis	
	☐ Primary immunodeficiency (eg, common variable immunodefici	
	combined immunodeficiency, Wiskott-Aldrich syndrome)	enej, 11 innieu ugummugiee uniiemu, se vere
	☐ Pediatric autoimmune neuropsychiatric disorders associated with	streptococcal infections (PANDAS)
	Pediatric acute onset neuropsychiatric syndrome (PANS)	i sucptococcai infections (i Ai (DAS)
	Other	
	Outer	
2.	What is the ICD-10 code? If patient's diagno	sis is PANDAS or PANS, no further questions
Co	mplete the following section based on the patient's diagnosis, if app	olicable.
Sec	ction A: Primary Immunodeficiency	
3.	Is this a request for continuation of immune globulin therapy? $\Box$	Yes ☐ No If No, skip to #8
4	The decree of the control of the con	1.1.6.4.1
4.	Has the patient experienced a reduction in the frequency of bacteria	al infections since starting immune globulin
	therapy? ☐ Yes ☐ No	
5.	Does the prescriber measure trough IgG levels at least once per year	r?
•	$\square$ Yes $\square$ No $\square$ Not applicable to diagnosis If Not applicable to	
	Tes Tho applicable to diagnosis if not applicable to	dugnosis, no juriner questions.
6.	Is the most recent trough IgG level at or above the lower range of n	
	ACTION REQUIRED: If 'yes', attach a copy of the laboratory rep	port with a recent IgG trough level.
	If Yes or Not applicable, no further questions. \(\sigma\) Yes \(\sigma\) No \(\sigma\)	Not applicable for diagnosis
_		
7.	Will the prescriber re-evaluate the dose of immune globulin and con	
	appropriate)? ☐ Yes ☐ No ☐ Not applicable/not clinically app	ropriate <i>ivo jurtner questions</i> .

8.	What is the specific immunodeficiency disorder?  ☐ Severe combined immunodeficiency (SCID), specify: ☐ Congenital agammaglobulinemia (eg, X-linked or autosomal recessive agammaglobulinemia) ☐ Other non-SCID combined immunodeficiency disorder, specify: ☐ Common variable immunodeficiency (CVID) ☐ Hypogammaglobulinemia (unspecified) or other predominant antibody deficiency disorder ☐ Selective IgA deficiency ☐ Selective IgM deficiency ☐ IgG subclass deficiency ☐ Specific antibody deficiency
	☐ Other immunodeficiency disorder/none of the above, <i>specify</i> :
9.	<ul> <li>ACTION REQUIRED: Please indicate and attach a copy of the following pre-treatment laboratory information (where applicable):</li> <li>IgG (total) level: mg/dL  a) Is IgG (total) level within the normal reference range? ☐ Yes ☐ No  b) If No, is the IgG level greater than or equal to (≥) 2 SD below the mean for age? ☐ Yes ☐ No</li> <li>IgG subclass levels:  a) IgG1 mg/dL; b) IgG2 mg/dL; c) IgG3 mg/dL;  d) Other  e) Are the IgG subclass levels within the normal reference range? ☐ Yes ☐ No  f) If No, is the level(s) greater than or equal to (≥) 2 SD below the mean for age? ☐ Yes ☐ No  g) Were IgG subclass levels measured on at least 2 different occasions? ☐ Yes ☐ No</li> <li>IgA level: mg/dL; Is the IgA level within the normal reference range? ☐ Yes ☐ No</li> <li>IgM level: mg/dL; Is the IgM level within the normal reference range? ☐ Yes ☐ No</li> </ul>
10.	If diagnosis is severe combined immunodeficiency, are maternal T cells present in the circulation? If Yes, skip to #12. $\square$ Yes $\square$ No
11.	If diagnosis is severe combined immunodeficiency, what is the patient's CD3 T cell count?  ACTION REQUIRED: Attach a copy of the laboratory report with lymphocyte subset enumeration by flow cytometry per microliter
12.	Was the diagnosis confirmed by molecular or genetic testing? <i>ACTION REQUIRED: Please attach a copy of the laboratory report or other medical record that shows the results of molecular/genetic testing.</i> Yes Do Do Not applicable to diagnosis
13.	If the diagnosis is common variable immunodeficiency, have other causes of immune deficiency been excluded (eg, drugs, infectious disease, malignancy)? $\square$ Yes $\square$ No $\square$ Not applicable to diagnosis
14.	Does the patient have a history of recurrent bacterial infections (eg, pneumonia, otitis media, sinusitis, sepsis, gastrointestinal infections)? ☐ Yes ☐ No
15.	Was the immune globulin therapy initiated in the hospital setting? ☐ Yes ☐ No
16.	Has the patient demonstrated an impaired antibody response to vaccination with a pneumococcal polysaccharide vaccine? <i>ACTION REQUIRED: If Yes, please attach a copy of the laboratory report with post-vaccination titers.</i> $\square$ Yes $\square$ No $\square$ Not applicable
	tion B: Chronic Inflammatory Demyelinating Polyneuropathy (CIDP)  Is this a request for continuation of immune globulin therapy? If Yes, skip to #21  Yes  No
18.	Is the disease course progressive or relapsing/remitting for 2 months or longer? ☐ Yes ☐ No
19.	Does the patient have moderate to severe functional disability? $\square$ Yes $\square$ No
20.	Were electrodiagnostic studies (electromyography [EMG] or nerve conduction studies [NCS]) and the evaluation of cerebrospinal fluid (when available) performed to confirm the diagnosis? <i>ACTION REQUIRED: If 'yes'</i> , <i>attach a copy of the EMG or NCS test results and CSF analysis.</i> $\square$ Yes $\square$ No <i>No further questions</i>

Send completed form to: Case Review Unit CVS Caremark Specialty Programs Fax: 1-855-330-1720

Note: This fax may contain medical information that is privileged and confidential and is solely for the use of individuals named above. If you are not the intended recipient you hereby are advised that any dissemination, distribution, or copying of this communication is prohibited. If you have received the fax in error, please immediately notify the sender by telephone and destroy the original fax message. Cuvitru PANDAS SOC SGM CFT – 03/2021.

CVS Caremark Specialty Pharmacy • 2211 Sanders Road NBT-6 • Northbrook, IL 60062

Phone: 1-888-877-0518 • Fax: 1-855-330-1720 • www.caremark.com

21.	Has the patient demonstrated significant improvement in disability and maintenance of improvement since starting IG therapy? $\square$ Yes $\square$ No
22.	Is IG being used at the lowest effective dose and frequency? ☐ Yes ☐ No
	tion C: Multifocal Motor Neuropathy (MMN)  Is this a request for continuation of immune globulin therapy? If Yes, skip to #26  Yes  No
24.	Has the patient experienced progressive, multifocal, asymmetrical weakness without objective sensory loss in 2 or more nerves for at least 1 month? $\square$ Yes $\square$ No
25.	Were electrodiagnostic studies (electromyography [EMG] or nerve conduction studies [NCS]) performed to confirm the diagnosis? <i>ACTION REQUIRED: If 'yes'</i> , <i>attach a copy of the EMG or NCS test results</i> . □ Yes □ No
26.	Has the patient demonstrated significant improvement in disability and/or maintenance of improvement since starting IG therapy? $\square$ Yes $\square$ No
	tion D: Dermatomyositis (DM) or Polymyositis (PM) Is this request for continuation of immune globulin therapy? If Yes, skip to #32
28.	Does the patient exhibit any of the following clinical features? <i>Indicate ALL that apply</i> .  ☐ Proximal muscle weakness (upper or lower extremity and trunk) ☐ Elevated serum creatine kinase (CK) or aldolase level ☐ Muscle pain on grasping or spontaneous pain ☐ Myogenic changes on EMG (short-duration, polyphasic motor unit potentials with spontaneous fibrillation potentials) ☐ Positive for anti-synthetase antibodies (e.g., anti-Jo-1, also called histadyl tRNA synthetase) ☐ Non-destructive arthritis or arthralgias ☐ Systemic inflammatory signs (fever: more than 37°C at axilla, elevated serum CRP level or accelerated ESR of more than 20 mm/h by the Westergren method ☐ Pathological findings compatible with inflammatory myositis (inflammatory infiltration of skeletal evidence of active regeneration may be seen) ☐ None of the above
29.	Were electrodiagnostic studies (electromyography [EMG]) and the muscle biopsy (when available) performed to confirm the diagnosis? <i>ACTION REQUIRED: If 'Yes', attach a copy of the EMG test results</i> $\square$ Yes $\square$ No
30.	Were standard first-line (corticosteroids) and second-line (immunosuppressants) treatments tried but were unsuccessful or not tolerated? <i>ACTION REQUIRED: If 'Yes', attach supporting chart note(s) describing previous treatments and no further questions.</i> $\square$ Yes $\square$ No
31.	Is the patient unable to receive standard first-line and second-line therapy because of a contraindication or other clinical reason? <i>ACTION REQUIRED: If 'Yes'</i> , attach supporting chart note(s) describing previous treatments. ☐ Yes ☐ No No further questions
32.	Has the patient demonstrated significant improvement in disability and/or maintenance of improvement since starting IG therapy? ☐ Yes ☐ No
	tion E: Parvovirus B19-Induced Pure Red Cell Aplasia (PRCA)  Does the patient have severe, refractory anemia associated with bone marrow suppression?   Yes  No
34.	Does the patient have parvovirus B19 viremia? ACTION REQUIRED: If 'yes', attach test result confirming presence of parvovirus B19. $\square$ Yes $\square$ No
	tion F: Myasthenia Gravis  What is the primary reason IG is being prescribed?  □ Refractory myasthenia gravis, <i>skip to #38</i> □ Acute exacerbation/crisis  □ Worsening weakness, <i>skip to #37</i>

	☐ Pre-operative management (eg, prior to thymectomy), <i>no further questions</i> ☐ Other
36.	Does the patient have severe swallowing difficulty and/or respiratory failure? <i>If Yes, no further questions.</i> □ Yes □ No
37.	Does the patient have weakness with an increase in any of the following symptoms: diplopia, ptosis, blurred vision difficulty speaking (dysarthria), difficulty swallowing (dysphagia), difficulty chewing, impaired respiratory status, fatigue, or limb weakness? $\square$ Yes $\square$ No <i>No further questions</i>
38.	Has the patient tried and failed 2 or more standard therapies (eg, corticosteroids, azathioprine, cyclosporine, mycophenolate mofetil, rituximab)? <i>ACTION REQUIRED: If 'Yes', attach supporting chart note(s) describing previous treatments.</i> □ Yes □ No
	tion G: Stiff-Person Syndrome  Has the diagnosis been confirmed by anti-glutamic acid decarboxylase (GAD) antibody testing?  ACTION REQUIRED: If 'Yes', attach GAD antibody test results.   Yes No
40.	Has the patient received first-line treatment with benzodiazepines and/or baclofen and experienced an inadequate response? <i>ACTION REQUIRED: If 'Yes'</i> , attach supporting chart note(s) describing previous treatments. ☐ Yes ☐ No
	tion H: Immune Thrombocytopenic Purpura (ITP)  Is the patient a pregnant woman? □ Yes □ No  If yes, please provide estimated date of delivery and no further questions:
42.	Is the patient an adult with refractory ITP after splenectomy? $\square$ Yes $\square$ No If No, skip to #45
43.	What is the current pre-treatment platelet count? ACTION REQUIRED: Attach lab report with platelet count per mcL If less than 30,000/mcL, no further questions.
44.	Does the patient have significant bleeding symptoms (eg, mucosal bleeding or other moderate to severe bleeding)? $\square$ Yes $\square$ No <i>No further questions</i>
45.	Is the patient at high risk for bleeding or does the patient require a rapid increase in platelets? If yes, please indicate the risk factors for bleeding or reason for a rapid increase in platelets.  Undergoing a medical or dental procedure where blood loss is anticipated Comorbidity (eg, peptic ulcer disease or hypertension) Mandated anticoagulation therapy Profession or lifestyle predisposes the patient to trauma (eg, construction worker, fireman, professional athlete) Other No, not at high risk or does not require rapid increase in platelets
46.	What is the current pre-treatment platelet count? <i>ACTION REQUIRED: Attach lab report with platelet count.</i> mcL
47.	Does the patient have significant bleeding symptoms (eg, mucosal bleeding or other moderate to severe bleeding)? $\square$ Yes $\square$ No
48.	What is the classification of ITP?  □ Newly-diagnosed ITP (diagnosed within the past 3 months), no further questions if patient is less than 18 years  □ Previously untreated ITP (initial therapy), no further questions if patient is less than 18 years old  □ Chronic or persistent ITP (greater than or equal to 3 months from diagnosis), skip to #51  □ ITP unresponsive to first-line treatment, skip to #51  □ Other
49.	Please indicate the prescribed regimen.  ☐ IG monotherapy ☐ IG in combination with corticosteroid, no further questions ☐ Other

50.	Is corticosteroid therapy contraindicated? $\square$ Yes $\square$ No No further questions
51.	Does the patient have relapsed ITP after a previous response to IG therapy? If Yes, no further questions. $\square$ Yes $\square$ No
52.	Does the patient have a history of inadequate response, intolerance or a contraindication to corticosteroid or anti-D therapy? <i>ACTION REQUIRED: If 'Yes'</i> , attach supporting chart note(s) describing previous treatments or contraindication. $\square$ Yes $\square$ No
Sec	tion I: B-Cell Chronic Lymphocytic Leukemia (CLL), Bone Marrow Transplant/Hematopoietic Stem Cell Transplant
	ipient Is this request for continuation of immune globulin therapy? If Yes, skip to #58 ☐ Yes ☐ No
54.	Is IG prescribed for prophylaxis of bacterial infections? ☐ Yes ☐ No
55.	What is the patient's pre-treatment IgG level? ACTION REQUIRED: If IgG is less than 500 mg/dL, attach a copy of the laboratory report with the pre-treatment IgG level mg/dL
56.	If diagnosis is B-cell chronic lymphocytic leukemia, does the patient have a history of recurrent sinopulmonary infections requiring intravenous antibiotics or hospitalization? $\square$ Yes $\square$ No No further questions
57.	If diagnosis is bone marrow transplant/hematopoietic stem cell transplant recipient, has the patient received a bone marrow/hematopoietic stem cell transplant within the past 100 days?   Yes No No further questions
58.	Has the patient experienced a reduction in the frequency of bacterial infections since starting IG therapy? $\square$ Yes $\square$ No
	tion J: HIV Infection: Prophylaxis or Thrombocytopenia  Is the requested drug being prescribed for prophylaxis of bacterial infections in a pediatric patient?  If Yes, skip to #70  Yes  No
60.	Is the requested drug being prescribed for treatment of thrombocytopenia associated with HIV? $\ \square$ Yes $\ \square$ No
61.	Is the patient an adult? ☐ Yes ☐ No If No, skip to #66
62.	Does the patient have significant bleeding? □ Yes □ No
63.	What is the patient's platelet count?/ mcL
64.	Is the patient Rh-positive? ☐ Yes ☐ No If No, no further questions.
65.	Has the patient failed treatment with RhIG? ☐ Yes ☐ No No further questions
66.	What is the patient's pre-treatment IgG level? ACTION REQUIRED: If IgG is less than 400 mg/dL, attach a copy of the laboratory report with the pre-treatment IgG levelmg/dL
67.	Has the patient had 2 or more bacterial infections in a 1-year period despite antibiotic chemoprophylaxis with TMP-SMZ or another active agent? If Yes, no further questions. $\square$ Yes $\square$ No
68.	Does the patient have HIV-associated thrombocytopenia despite anti-retroviral therapy? <i>If Yes, no further questions.</i> $\square$ Yes $\square$ No
69.	What is the patient's T4 cell count? / mm3
70.	Is this request for continuation of immune globulin therapy? If Yes, skip to #79 ☐ Yes ☐ No
71.	Please indicate whether IG will be used for primary or secondary prophylaxis.  □ Primary prophylaxis □ Secondary prophylaxis, <i>skip to #73</i> □ Other, <i>skip to #74</i>
72.	What is the patient's pre-treatment IgG level? ACTION REQUIRED: If IgG is less than 400 mg/dL, attach a copy of the laboratory report with the pre-treatment IgG level mg/dL  If less than 400 mg/dL, no further questions. If greater than or equal to 400 mg/dL, skip to #77

73.	Does the patient have a history of recurrent bacterial infections (greater than 2 serious bacterial infections in a 1-year period)? If Yes, no further questions. $\square$ Yes $\square$ No
74.	Has the patient failed to form antibodies to common antigens, such as measles, pneumococcal, and/or Haemophilus influenzae type b vaccine? <i>If Yes, no further questions.</i> □ Yes □ No
75.	Is this request for a single dose of immune globulin for a patient who has been exposed to measles? If Yes, no further questions. $\square$ Yes $\square$ No
76.	Does the patient live in an area where measles is highly prevalent? $\square$ Yes $\square$ No If No, skip to #78
77.	Has the patient failed to develop an antibody response after two doses of measles, mumps, and rubella live virus vaccine? If Yes, no further questions. $\square$ Yes $\square$ No
78.	Does the patient have chronic bronchiectasis that is suboptimally responsive to antimicrobial and pulmonary therapy? $\square$ Yes $\square$ No <i>No further questions</i>
79.	Has the patient experienced a reduction in the frequency of bacterial infections since starting IG therapy? ☐ Yes ☐ No
	tion K: Lambert-Eaton Myasthenic Syndrome
80.	Is this request for continuation of immune globulin therapy? If Yes, skip to #86 ☐ Yes ☐ No
81.	Has the diagnosis been confirmed by neurophysiology studies (e.g., electromyography) or a positive anti- P/Q type voltage-gated calcium channel antibody test? <i>ACTION REQUIRED: If 'yes', attach a copy of the laboratory report, neurophysiology study report or other supporting medical record(s).</i> ☐ Yes − Neurophysiology studies
	☐ Yes – Positive anti- P/Q type voltage-gated calcium channel antibody test☐ No
82.	Has the patient tried an anticholinesterase (e.g., pyridostigmine) but it was unsuccessful or not tolerated? ☐ Yes ☐ No
83.	Has the patient tried amifampridine (eg 3,4-diaminopyridine phosphate, Firdapse) but it was unsuccessful or not tolerated? ☐ Yes ☐ No
84.	Does the patient have severe weakness? If Yes, no further questions. □ Yes □ No
85.	Is there difficulty with venous access for plasmapheresis? $\square$ Yes $\square$ No No further questions
86.	Has the patient experienced stability or improvement in symptoms relative to the natural course of LEMS? ☐ Yes ☐ No
	tion L: Immune Checkpoint Inhibitor-Related Adverse Events
87.	Has the patient experienced a moderate or severe adverse event to a PD-1 inhibitor (e.g., pembrolizumab, nivolumab) or PD-L1 inhibitor (e.g., atezolizumab, avelumab, durvalumab)? ☐ Yes ☐ No
88.	Is the offending drug being temporarily held or has it been discontinued permanently? $\ \square$ Yes $\ \square$ No
89.	Which of the following adverse events did the patient experience?  ☐ Pneumonitis ☐ Peripheral neuropathy ☐ Myasthenia gravis ☐ Encephalitis ☐ Other
	tion M: Hypogammaglobulinemia from CAR-T Therapy  Has the patient received treatment with CAR-T therapy (e.g., tisagenlecleucel [Kymriah] or axicabtagene ciloleuce [Yescarta]?
91.	What is the patient's IgG level? ACTION REQUIRED: If IgG is less than 400 mg/dL, attach a copy of the laboratory report with the pre-treatment IgG level mg/dL  Unknown
Sec	tion N: Guillain-Barre Syndrome (GBS)

92. Does the patient have severe disease with significant weakness (e.g., inability to stand or walk without aid, respiratory weakness)?   Yes No	
93. Did the onset of neurologic symptoms occur less than 4 weeks from the anticipated start of immunoglobulin therapy? $\square$ Yes $\square$ No	
Section O: Acute Disseminated Encephalomyelitis  94. Has the patient had an insufficient response to intravenous corticosteroid treatment?   Yes   No	
Section P: Autoimmune Mucocutaneous Blistering Disease (includes Pemphigus Vulgaris, Pemphigus Foliaceus, Bullous Pemphigoid, Mucous Membrane Pemphigoid, and Epidermolysis Bullosa Aquisita)  95. Has the diagnosis been proven by biopsy and confirmed by pathology report? ☐ Yes ☐ No	-
96. Is the condition rapidly progressing, extensive, or debilitating? ☐ Yes ☐ No	
97. Has the patient failed or experienced significant complications (eg diabetes, steroid-induced osteoporosis) frostandard treatment (corticosteroids, immunosuppressive agents)?   Yes  No	om
Section Q: Autoimmune Hemolytic Anemia  98. Which type of autoimmune hemolytic anemia does the patient have?  □ Warm type □ Cold type □ Other	
99. Has the patient tried corticosteroids with inadequate response? If Yes, no further questions. $\square$ Yes $\square$ No	
100. Has the patient has a splenectomy with inadequate response? <i>If Yes, no further questions.</i> □ Yes □ No	
101.Does the patient have a contraindication to corticosteroids or splenectomy? ☐ Yes ☐ No	
Section R: Autoimmune Neutropenia 102.Is treatment with G-CSF (granulocyte colony stimulating factor) an appropriate option? Examples of G-CSF Fulphila, Granix, Leukine, Neulasta, Neuopogen, Udenyca, Zarxio. □ Yes □ No	include
Section S: Birdshot Retinochoroidopathy 103.Has the patient tried immunosuppressant therapy (e.g., corticosteroids, cyclosporine) with inadequate respon ☐ Yes ☐ No	se?
Section T: Churg-Strauss Syndrome 104.Does the patient have severe, active disease? □ Yes □ No	
105. Will immune globulin be used as adjunctive therapy? ☐ Yes ☐ No	
106.Has the patient experienced failure, intolerance, or is contraindicated to other interventions? $\square$ Yes $\square$ No	
Section U: Enteroviral Meningoencephalitis 107.Is the patient's condition severe? □ Yes □ No	
Section V: Hematophagocytic Lymphohistiocytosis (HLH) and Macrophage Activation Syndrome (MAS) 108. What is the patient's total IgG level? ACTION REQUIRED: Attach a copy of the laboratory report with the treatment IgG level mg/dL If less than 400 mg/dL, no further questions.	e pre-
109.Is the total IgG level at least two standard deviations below the mean for age? ☐ Yes ☐ No	
Section W: Hyperimmunoglobulinemia E syndrome 110.Does the patient have severe eczema? □ Yes □ No	
Section X: Multiple Myeloma 111.Does the patient have recurrent, serious infections despite the use of prophylactic antibiotics? □ Yes □ N	o
Section Y: Neonatal Hemochromatosis 112.Is the patient currently pregnant? □ Yes □ No	
113.Does the patient have a history of pregnancy ending in documented neonatal hemochromatosis? $\Box$ Yes	No
Section Z: Opsoclonus-Myoclonus	

Prescriber or Authorized Signature Date (mm/dd/yy)		
X		
I attest that this information is accurate and true, and that documentation supporting this information is available for review if requested by CVS Caremark or the benefit plan sponsor.		
131. Does the patient have persistent oliguria with pulmonary edema? ☐ Yes ☐ No		
130. Does the patient have an undrainable focus of infection? If Yes, no further questions. $\square$ Yes	<b>l</b> No	
129. Is the infection refractory to several hours of aggressive therapy? If Yes, no further questions. $\Box$	Yes No	
Section GG: Toxic Shock Syndrome  128.Does the patient have toxic shock syndrome due to a staphylococcal or streptococcal infection?  **ACTION REQUIRED: If 'yes', attach culture or Gram stain. □ Yes □ No		
Section FF: Toxic Necrotizing Fasciitis  127. Does the patient have toxic necrotizing fasciitis due to invasive group A streptococcal infection?  **ACTION REQUIRED: If 'yes', attach documentation confirming presence of fasciitis and cultustain. □ Yes □ No	ure or Gram	
126. Has the patient experienced inadequate response, intolerance, or have a contraindication to second ☐ Yes ☐ No	l line therapy?	
125. Has the patient experienced inadequate response, intolerance, or have a contraindication to first lin ☐ Yes ☐ No	ne therapy?	
Section EE: Systemic Lupus Erythematosus 124.Does the patient have severe, active disease? ☐ Yes ☐ No		
Section DD: Toxic Epidermal Necrolysis, Stevens-Johnson Syndrome 123. Is the patient's case severe? □ Yes □ No		
122. What is the patient's pre-treatment IgG level? ACTION REQUIRED: If IgG is less than 400 mg/of the laboratory report with the pre-treatment IgG level mg/dL  Unknown	/dL, attach a copy	
Section CC: Secondary Immunosuppression Due to Surgery, Malignancy, Burns, Collagen-Vascular D 121.Is immune globulin being requested to prevent or modify recurrent bacterial or viral infections?		
120.Is the patient undergoing renal transplantation from a live donor with ABO incompatibility or posi ☐ Yes ☐ No	itive cross match?	
Section BB: Solid Organ Transplantation 119. Is immune globulin being prescribed for solid organ transplantation in an allosensitized patient?  If Yes, no further questions. □ Yes □ No		
118. Did the patient try corticosteroids with no improvement in symptoms? $\square$ Yes $\square$ No		
Section AA: Rasmussen Encephalitis 117.Did the patient try anti-epileptic drugs with no improvement in symptoms? ☐ Yes ☐ No		
116.Is immune globulin being used as last-resort treatment? ☐ Yes ☐ No		
115.Does the patient have refractory opsoclonus-myoclonus? ☐ Yes ☐ No		
114. Does the patient have paraneoplastic opsoclonus-myoclonus-ataxia associated with neuroblastoma <i>If Yes, no further questions.</i> □ Yes □ No	n?	